


# CYCLE

## Connectivity of Coral Ecosystems in the northern Gulf of Mexico

Annalisa Bracco






**Santiago Herrera**  
**LeHigh University**




**Andrea Quattrini**  
**Harvey Mudd College**



**Annalisa Bracco**  
**Georgia Tech**



**Randy Clark**  
**NOAA**



**Andrew David**  
**NOAA**

**cycle**



**Peter Etnoyer**  
**NOAA**



**Amanda Demopoulos**  
**USGS**




**Stacey Harter**  
**NOAA**



**Christopher Meyer**  
**Smithsonian**



**Nancy Prouty**  
**USGS**




**Joshua Voss**  
**Harbor Branch**  
**Ocean. Inst.**

# PROBLEM

- Continued anthropogenic threats in the marine environment → urgent need to to effective **manage and conserve vulnerable coral ecosystems** in the Gulf of Mexico (GoM).
- **Marine protected areas (MPAs)** are essential to protect and conserve coral ecosystems → **key restoration strategy for benthic communities** impacted by the *Deepwater Horizon* oil spill.
- The **Flower Garden Banks National Marine Sanctuary (FGBNMS)** has proposed to **expand current boundaries** to encompass additional coral sites.



# Flower Garden Banks National Marine Sanctuary Expansion: Sanctuary Advisory Council Recommendation May 2018




# CYCLE

A **collaborative effort** to enhance understanding of GoM ecosystems and link research with regional conservation initiatives to 1) **inform the restoration of degraded deepwater coral communities**, and 2) **preserve long-term viability of coastal ecosystems**.







CYCLE will

- provide ecosystem connectivity information and tools to managers to effectively manage the FGBNMS MPAs, the boundary expansion alternatives, and areas further afield.
- improve understanding of **the processes that shape 3D connectivity networks in shallow (15-40m), upper mesophotic (40-85m), lower mesophotic (85-150m) and deepwater (> 150m) coral ecosystems in the GoM.**

- Focus on twelve species: two hard corals (*Montastraea cavernosa* and *Orbicella faveolata*), two soft corals (*Swiftia exserta* and *Hypnogorgia pendula*), one black coral (*Stichopathes lutkeni*), one sponge (*Xestospongia muta*), two fishes (red snapper *Lutjanus campechanus* and tomtate *Haemulon aurolineatum*) and four octocoral species (*Hypnogorgia pendula* and *Swiftia exserta* from mesophotic areas; *Callogorgia delta* from the upper continental slope and *Paramuricea biscaya*).
- Integration of field sampling and genetic, chemical, and ecological analyses with habitat suitability, oceanographic, and larval dispersal modeling.
- Additionally, analysis of coral recruitment and species biodiversity through the use of Autonomous Reef Monitoring Structures or ARMS.





# How



**CYCLE**



# Potential connectivity of *Callogorgia delta* in the Northern Gulf of Mexico





# Testing processes involved in the depth differentiation hypothesis

The bathyal region (~200 – 2000 m) is a source of genetic diversity + a high rate of species formation → genetic differentiation should occur over relatively small vertical distances

Is it true in the northern Gulf of Mexico?

What is the controlling factor?







# Genetic structuring with distance



*Callogorgia delta*




61,179 loci

Fst 0.0115-0.0691



# PATTERN FORMATION AND SUBMESOSCALES

## Transport along the continental slope




Bracco, Choi et al. Ocean Modelling, 2016

# GENERATION OF SUBMESOSCALE EDDIES ALONG CONTINENTAL SLOPES


- **Vertical velocities have to go to zero at the bottom -> horizontal shear layer near the bottom**
- **Juxtaposition of along-slope frontal currents that are highly variable in speed and direction (very common in the Gulf of Mexico) -> lateral shear layers**
- The width of the layer depends on bathymetric slope; mostly unresolved at  $\geq 5$ km horizontal resolution and partially resolved at  $\sim 1$  km
- Whenever high values of vorticity are achieved ( $\zeta/f > 0.4$ ) small scales eddies and filaments are generated through partially unbalanced instabilities

# Juxtaposition of along-slope frontal currents that are highly variable in speed and direction -> lateral shear layers


mean  $u$  (cm/s)




instantaneous  $u$  (cm/s)



$\zeta/f$




Vertical velocities have to be zero at the bottom -> horizontal shear layer near the bottom




instantaneous  $u$  (cm/s)

$\zeta/f$




**Observed (dashed lines;** Jim Ledwell deep release experiment; Ledwell et al., 2017) and **modeled (red)** diapycnal diffusivity in the northern GoM



Snapshot of relative vorticity in the bottom layer 10<sup>th</sup>, 2015

# Virtual connectivity using a regional circulation model









*Callogorgia delta*


PLD = 40 Days

**ROMS**

1 km horizontal resolution  
4 larvae/particle releases in each season in 2015 and 2016



# Modeled connectivity



the limited connectivity is controlled by diapycnal mixing



# Conclusions

- The new analysis confirms the presence of a pattern of genetic differentiation in *C. delta* across a gradient of depth
- Abiotic gradients associated influence the evolution of deep-sea populations and species
- Future design of protected areas and the Flower Garden Banks expansion should incorporate a variety of depth ranges to capture the diversity within and among vulnerable marine ecosystems

